

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
12 July 2001 (12.07.2001)

PCT

(10) International Publication Number
WO 01/49257 A1

(51) International Patent Classification⁷: **A61K 7/48, 7/06**

(21) International Application Number: PCT/US00/35469

(22) International Filing Date:

28 December 2000 (28.12.2000)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

09/478,071

1 January 2000 (01.01.2000)

US

(71) Applicant: **INTERNATIONAL FLORA TECHNOLOGIES, LTD.** [US/US]; 1151 North Fiesta Blvd., Gilbert, AZ 85233-2238 (US).

(72) Inventors: **BROWN, James, H.**; 5214 East Paradise Drive, Scottsdale, AZ 85224 (US). **COPELAND, Lee, Roy**; 8716 E. Edgewood, Mesa, AZ 85208 (US). **KLEIMAN, Robert**; 7129 East Laguna Azul Avenue,

Mesa, AZ 85208 (US). **KORITALA, Sambasivarao**; 1104 West Stacey Lane, Tempe, AZ 85284 (US). **CUMMINGS, Melanie, K.**; 4063 West Orchid Lane, Chandler, AZ 85226 (US).

(74) Agent: **HALVORSON, Kristofer, E.**; The Halvorson Law Firms P.C., Ste. #1, 405 W. Southern Ave., Tempe, AZ 85282 (US).

(81) Designated States (*national*): CA, JP.

(84) Designated States (*regional*): European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR).

Published:

- With international search report.
- With amended claims.

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: HIGH UNSAPONIFIABLES AND METHODS OF USING THE SAME

(57) **Abstract:** Materials with high levels of unsaponifiable matter result in Hydrolysates with unique properties. It has been found that the application of a hydrolysis process to materials, particularly materials with a high level of unsaponifiables (e.g. at least 6% by total weight of the material) produces a product with properties significantly different from those products resulting from the conventional saponification of materials with less than 6% by weight of unsaponifiables. The resulting Hydrolysates resist both physical and aqueous-based removal from skin and hair, exhibit a very unique surfactant property, and are not foaming agents with water. Hydrolysates according to the present invention may thus be used to enhance the performance of cosmetics and pharmaceuticals. These Hydrolysates are bioactive agents and alternative natural carrying agents for topical application of materials, particularly for application of materials to the skin or hair of subjects, and provide a substantive support for the materials carried.



WO 01/49257 A1

**HIGH UNSAPONIFIABLES AND METHODS OF USING THE
TECHNICAL FIELD OF THE INVENTION**

The present invention relates to a novel composition of matter derived from natural materials or extracts of natural materials. In particular the invention relates to substantive carriers derived from natural waxes, oils, and extracts, and in particular to substantive carriers derived from natural ingredients with relatively high levels of unsaponifiable materials (as defined below) and methods of using the same.

BACKGROUND OF THE ART

Vegetable and animal fats are organic lipid materials that generally contain esters of long-chain fatty acids and glycerine. Under certain conditions these esters react with water (hydrolysis) form an alcohol (glycerine) and fatty acids. (Hydrolysis is the splitting of a compound into components by the addition of water and an enzyme, acid or base.) The results of a hydrolysis reaction are known as "Hydrolysates". When heated in the presence of an alkali hydroxide the above mentioned esters yield soap (the alkali salt of the fatty acid) and glycerine; this particular hydrolysis process is called saponification. "Saponification" and "saponifying" are used herein in their normal manner to mean the hydrolysis reaction between a wax, oil or fat with an alkali metal or alkaline earth metal hydroxide to form the corresponding metallic salt soap. These fats and oils have a saponification value that is the number of milligrams of potassium hydroxide required for complete saponification of one gram of free organic acid and/or organic acid ester.

The post saponification products may either be hydrophilic (water soluble) or hydrophobic (water insoluble). Herein we will use the term "unsaponifiable" to mean those materials that, after the saponification reaction is completed, remain water insoluble. This is in full accord with the A.O.C.S. Official Method Ca 6b-53, which defines unsaponifiable materials as those substances frequently found as components of fats and oils, which cannot be saponified by the usual caustic treatment, but which are soluble in ordinary fats and oils. Included, but not limiting, in the group of unsaponifiable materials are higher aliphatic alcohols, sterols, pigments, mineral oils, and hydrocarbons. Unsaponifiable materials are generally non-volatile at 103°C. The weight percent of unsaponifiable material in a substance may be measured directly by measuring the weight percent of those materials defined as unsaponifiable.

The most well known vegetable and animal lipids have low levels, less than five percent (<5%), of unsaponifiable materials. This means that most of the products of the saponification reaction are water-soluble. Commonly used vegetable oils have levels of unsaponifiable materials generally below 1%. For example, saponification of soybean oil leaves 0.7 weight percent

unsaponifiable materials, saponification of olive oil leaves 1.2 weight percent unsaponifiable materials, and saponification of peanut oil leaves 0.4 weight percent unsaponifiable materials. However, some commercial oils contain higher concentrations of unsaponifiable products, up to as much as 6.0 weight percent unsaponifiable materials. Examples include: crude rice bran oil, 4.2 %
5 unsaponifiables, crude wheat germ oil, 6 % unsaponifiables, and shea butter, 9-13 % unsaponifiables. Materials with high levels of unsaponifiables, such as shea butter, are not a preferred starting material for the production of soap because of the high amount of unsaponifiable materials left after the saponification reaction.

In most cases, the hydrolysis products of a saponification process are used solely for a single
10 purpose, which is as a hygienic skin-cleansing agent (soap). In the past, the basic ingredient of soap was animal fat (also known as lard or tallow) with wood ash based lye used in the saponification process. Ideally a bar of soap has a suitable hardness to maximize user cycles and has a certain amount of resistance to water reabsorption when not in use, while at the same time providing sufficient lather (acting as a foaming agent) to enhance the cleaning ability of the soap. Animal
15 lipids as the active ingredient in the soap making process will generally meet these user demands to a greater or lesser degree. Current soap production continues to rely heavily on animal fats in their products to meet consumer demand and production requirements, although more and different types of synthetic materials are beginning to find use in soap compositions. The various synthetic compounds and mixtures of compounds have become very popular additions in modern soap making
20 technology for their improvement to soap quality and user satisfaction. However, these synthetic based soaps are generally resistant to the natural breakdown processes (i.e. biodegradability) and are thus relatively persistent in the environment.

There are basically two distinct types of soap manufacturing processes. In a first method, oils and fats are boiled in an open kettle with caustic alkali solutions, bringing about saponification
25 gradually until all of the fats and oils are completely saponified, followed by the removal of the glycerine. This process may either run in batch or in a continuous process.

In a second method, which is typically a continuous method (but may be run in batch form), fatty acids and alkali are brought together in proper portions for complete saponification in a mixing valve or other device which brings them in intimate contact. The progress of saponification depends
30 on the temperature, time of contact and efficiency of mixing. Concentrated solutions produced by these methods are referred to as "neat" soaps, and possess a concentration of 60-65% soap, about 35% water and traces of salt and glycerine. It is from this product that consumer soaps in the form of bars, flakes, granules and powders are produced, by first drying the neat soap into pellets having a moisture content of about 12-16% followed by finishing steps, such as milling, plodding,

amalgamating, and the like.

Consumer bar soaps today are manufactured from coconut oil and/or tallow or their fatty acids. Palm kernel oil is sometimes substituted for coconut oil for economic reasons, and soaps prepared with palm kernel oil are adjusted for performance characteristics similar to non-substituted tallow/coconut formulations. Palm oil is also often substituted for tallow.

A consideration in selecting materials for making soap is the proper ratio of saturated versus unsaturated, and long-versus-short-chain fatty acids that result in a soap having the desired qualities of stability, solubility, ease of lathering, hardness, cleaning ability, and the like. It has been determined that soaps prepared from fatty acid mixtures wherein a majority of the fatty acids in the mixtures has carbon chains less than twelve atoms irritate skin. Soaps prepared from saturated C₁₆ and C₁₈ fatty acids are typically too insoluble for consumer use. Thus, the preferred materials for soap production have fatty acid chains between twelve and eighteen carbon atoms in length.

Saponification of tallow produces a soap comprised of a mixture of fatty acids of C_{14:0}, C_{16:0}, C_{18:0}, and C_{18:1} (myristic, palmitic, stearic and oleic acids, respectively) and saponification of coconut oil produces a soap comprised of a mixture of fatty acids of C_{12:0} and C_{14:0} (lauric acid and myristic acid, respectively) and significant amounts of C_{8:0} and C_{10:0} fatty acids. Consumer soap preparations usually contain tallow/coconut (T/C) ratio ranges from approximately 90:10 to 75:25. Since lauric acid is found only in the coconut fraction of T/C mixtures, the most dramatic change observed in increasing the percent of the coconut fraction of T/C mixtures is the increase in the lauric acid. Increasing the coconut fraction in T/C fatty acid containing soaps generally improves the desirable foaming characteristics of such soaps. However, in soaps with T/C ratios of 50:50, the desirable skin mildness properties are reduced.

Typical fatty acid distribution (in weight percent) of the main soap making components is given below:

Carbon Chain Length	Tallow	Palm	Coconut	Palm Kernel
10:0 (capric)	0.1	0.0	15.1	6.4
12:0 (lauric)	0.1	0.3	48.0	46.7
14:0 (myristic)	2.8	1.3	17.5	16.2
16:0 (palmitic)	24.9	47.0	9.0	8.6
18:0 (stearic)	20.4	4.5	9.0	8.6
18:1 (oleic)	43.6	36.1	5.7	16.1
18:2 (linoleic)	4.7	9.9	2.6	2.9
18:3 (linolenic)	1.4	0.2	0.0	0.0

20:0 (arachidic)	1.8	0.3	0.0	0.4
------------------	-----	-----	-----	-----

From the table it can be seen that the coconut and palm kernel fats (both known as the lauric fats) are particularly rich in the C₁₀₋₁₄ saturated fatty acids, particularly derivatives from lauric acid itself. Another fat that contains saturated, relatively short chain fatty acids similar to coconut oil is babassu oil. In contrast, tallow and palm oil *per se* are industrial sources of non-lauric fats, especially those containing C₁₆ and C₁₈ fatty acids.

In general the longer chain fatty acid alkali salts, particularly the less expensive C₁₆ and C₁₈ salts (as obtained from tallow and palm oils), provide structure in the finished soap bars and prevent or retard disintegration of the soap bar on exposure to water. The more expensive, shorter chain, lauric fat-derived, (i.e., lauric acid salts) and other soluble salts (typically as obtained from coconut and palm kernel oil) contribute to the lathering properties of the overall composition. A general problem in the formulation of bar soaps has been finding a balance between providing structure (generally obtained from the long chain component) and maintaining lathering properties (generally obtained from the more expensive short chain component) at a practical overall cost.

In addition to fatty acid salts, soap bars can contain free fatty acids. The addition of free fatty acids is known as 'superfatting'. Superfatting at a 5-10% free fatty acids level is known to give a copious, creamy lather. Other superfatting agents used include citric and other acids that function by promoting the formation of free fatty acids in the fat blend.

For the manufacture of the soap cakes, common additives can be added to the base soap in the normal quantities, referred to 100 parts by weight of base soap, such as overgreasing agents (1 to 3 wt.%), stabilizers (antioxidants, complexing agents) (0.05 to 0.5 wt.%), perfume (0.5 to 3 wt.%) and possibly dyes (0.05 to 0.3 wt.%) as well as skin protection agents such as sorbitol, glycerine or the like (1 to 5 wt.%).

The pharmaceutical and cosmetic industries have been using fat extracts of vegetable origin since earliest times. A number of years ago it became apparent in these industries that particularly valuable biological properties resulted from the use of vegetable fats or extracts of vegetable fats rich in unsaponifiable materials. Certain vegetable oils, for example avocado, and, in particular, shea butter, are known to be particularly rich in unsaponifiable materials and/or to contain, these unsaponifiable materials.

A process for enriching unsaponifiables in oils, especially shea butter, for use in cosmetic and pharmaceutical compositions is described in U.S. Pat. No. 5,679,393, issued to Laur. This process concentrates the unsaponifiable fraction of fats and oils by the processes of crystallization and fractionation. This method is expensive and it does not liberate the alcohol moiety from the

starting compounds (hydrolysis). Thus, the Laur process and methods for use of the products thereof never utilize hydrolysis to create alkali salts and liberate alcohols and other unsaponifiables.

Hydrolysates applied topically to animate and inanimate objects find use in numerous non-cleansing areas ranging from cosmetic preparations, pharmaceuticals, hydration formulations, insecticides, insect repellant, and the like. One of the areas of interest created by the varied uses of topically applied agents is maximizing the duration a topically applied active agent is present on the applied surface (substantivity). As a result of this intense interest, the search for ways to improve the duration of a fixed amount of topically applied cosmetics, pharmaceuticals, and bioactive agents has been of prime importance in all areas wherein topically applied cosmetics, pharmaceuticals, and bioactive agents are employed. An example of this interest may be found in the prior art relating to sunscreen compositions.

The use of sunscreen compositions is required by a large segment of society since only a small portion of those exposed to sunlight have the natural pigmentation which provides protection against the harmful effects of solar radiation. Because many people show erythema under even short exposures to sunlight, there is a need for sunscreen compositions that protect against erythema-causing radiation, i.e., ultraviolet radiation, so that longer exposure to the sunlight with less risk of sunburn is possible.

A variety of sunscreen compositions are known in the art. One tendency in formulating sunscreen compositions has been to prepare compositions that are water-resistant to the skin. One method is to chemically modify the ultraviolet absorber to increase its interaction with the skin by quaternizing imidazoles, as described in U.S. Pat. No. 3,506,758; another method is to copolymerize ultraviolet light absorbing monomers with other monomers to form water-resistant films, as described in U.S. Pat. Nos. 3,529,055 and 3,864,473; yet another method is to form polymeric films with water-insoluble polymers, as described in U.S. Pat. No. 3,784,488.

The use of the acid form of crosslinked ethylene-maleic anhydride copolymers to retain ultraviolet light absorbers is disclosed in U.S. Pat. No. 3,821,363. The use of water insoluble acrylate polymer having a solubility parameter of 6 to 10 in weak hydrogen bonding solvents is disclosed in U.S. Pat. No. 4,172,122. The use of water-insoluble, alcohol-soluble, film-forming poly-amide materials is disclosed in U.S. Pat. No. 3,895,104 solely for the purpose of providing improved substantivity.

The cosmetics and other applications of the prior art have not heretofore utilized the substantivity inherent in Hydrolysates of naturally derived materials containing high unsaponifiables or long chain esters (greater than 18 carbons in length) to enhance the intrinsic substantivity of topically applied agents with which they are incorporated. Previously, the purpose of employing

polymers or polymeric materials in the compositions of the prior art has been directed towards improving the adherency, i.e., substantivity, of the topical material to the skin or have been employed solely as thickening agents. The improved substantivity, among other properties, achieved by employing the Hydrolysates according to the present invention has not heretofore been disclosed or appreciated in the prior art.

The increased substantivity of topically applied agents provides for more effective and economical use of such materials. In particular, the present invention provides improved compositions, including emollients, skin hydrating agents, sunscreens, lipsticks, make up, insect repellants, insecticides, pesticides, herbicides, and the like, having at least an effective amount of a Hydrolysate including high levels of unsaponifiable materials, preferably of long chain esters.

SUMMARY OF THE INVENTION

The hydrolysis of materials with high levels of unsaponifiable matter, such as extracts from plants, result in products with unique properties. Traditional products of saponification of natural oils function as they do as a direct result of the low level of unsaponifiable contained therein (as discussed above). Such properties include high levels of aqueous surfactant activity, water-solubility or ready water-dispersability, activity as foaming agents, and the like. The very objective of traditional saponification processes is to increase the water-solubility and surfactant activity of naturally occurring materials. It has been found that the application of hydrolysis to materials, particularly naturally derived materials, with a high unsaponifiables fraction (e.g., at least 6% by total weight of the material) in combination with a saponifiable fraction produces a Hydrolysate with properties that are significantly different from those products resulting from the conventional saponification of materials with less than 6% by weight of unsaponifiable.

The resulting products from the practice of the present invention are substantive, water resistant, prevent unwanted absorption of a carried active ingredient by the applied surface, exhibit a unique surfactant functionality, and are not foaming agents with water. Some unexpected uses for the resulting Hydrolysates have been found to be as an emollient and/or an alternative natural carrying agent for topical application of cosmetics, pharmaceuticals, and bioactive agents, particularly to the skin of subjects, and provide a substantive support for the materials carried.

The novel features that are considered characteristic of the invention are set forth with particularity in the appended claims. The invention itself, however, both as to its structure and its operation together with the additional object and advantages thereof will best be understood from the following description of the preferred embodiment of the present invention when read in conjunction with the accompanying drawings. Unless specifically noted, it is intended that the words and phrases in the specification and claims be given the ordinary and accustomed meaning to those

of ordinary skill in the applicable art or arts. If any other meaning is intended, the specification will specifically state that a special meaning is being applied to a word or phrase. Likewise, the use of the words "function" or "means" in the Detailed Description of the Invention is not intended to indicate a desire to invoke the special provision of 35 U.S.C. §112, paragraph 6 to define the invention. To the contrary, if the provisions of 35 U.S.C. §112, paragraph 6, are sought to be invoked to define the invention(s), the claims will specifically state the phrases "means for" or "step for" and a function, without also reciting in such phrases any structure, material, or act in support of the function. Even when the claims recite a "means for" or "step for" performing a function, if they also recite any structure, material or acts in support of that means of step, then the intention is not to invoke the provisions of 35 U.S.C. §112, paragraph 6. Moreover, even if the provisions of 35 U.S.C. §112, paragraph 6, are invoked to define the inventions, it is intended that the inventions not be limited only to the specific structure, material or acts that are described in the preferred embodiments, but in addition, include any and all structures, materials or acts that perform the claimed function, along with any and all known or later-developed equivalent structures, materials or acts for performing the claimed function.

DESCRIPTION OF THE DRAWINGS

Figure 1 is a graph of the effects of the Hydrolysate when used with a hydrating lotion.

Figure 2 is a graph of the effects of the Hydrolysate when used with make up.

Figure 3 is an illustration of the average percentage of fly reduction when fly spray incorporates the Hydrolysates according to the present invention.

Figure 4 is the graph of the fly reduction for Aquapel when incorporating the Hydrolysates of the present invention.

Figure 5 is the graph of the fly reduction for Ceratex when incorporating the Hydrolysates of the present invention.

Figure 6 is the graph of the fly reduction for GnatAway when incorporating the Hydrolysates of the present invention.

Figure 7 is the graph of the fly reduction for Solitude when incorporating the Hydrolysates of the present invention.

DISCLOSURE OF INVENTION AND BEST MODE FOR CARRYING OUT INVENTION

The present invention is a composition of matter, and method for using the same, which is useful as a topically applied material with several useful inherent properties, such as substantivity. Additionally, the composition is useful for carrying an effective amount of topically applied active materials. More specifically, the composition according the present invention provides a carrying agent for the topical application of materials when superior "lasting" power or substantivity is

required. Additionally, the present invention is useful because, among other things, it acts as both an emollient and unique emulsifier and demonstrates substantivity; it has the ability to “fix” many different types of “active” materials, from sunscreens to pharmaceutical preparations to any applied animate or inanimate surface.

5 For the purposes of this invention, the following definitions should be considered:

“High unsaponifiable materials” or “high unsaponifiable content” oils, waxes, fats, and the like, means compositions that comprises at least 6 % by weight of total organic materials that are unsaponifiable and at least 10 % by weight of organic materials that are saponifiable (it is possible that the percentage of unsaponifiables may even exceed 95% in some formulations). Therefore, the
 10 term includes compositions containing from 6-90% by weight of organics of unsaponifiable materials and 10-94% by weight of saponifiable materials. Examples of bio-based materials with high unsaponifiables are listed in the table below.

Material	% Unsaponifiables
amaranth seed oil	9%
anise seed oil	7%
avocado seed oil	57%
barley oil	6%
briza oil	78%
buck wheat oil	7%
candelilla wax	65-75%
carnuba wax	50-55%
cassia occidentalis oil (wild coffee)	7%
coffee bean oil	8%
deoiled lecithin	32% (in Theory)
dog fish oil	16-18%
esparto wax	42-49%
oils from fungi and other microorganisms	6% or greater
guayule (plant material extract)	8-12%

jojoba oil	45%
jurinea oil	40%
lanolin	39%
laurel berry oil	6%
olestra(TM)or olean(TM)	33% (approximation)
olive oil concentrate (phytosqualene)	35-75%
olive seed oil	greater than 6%
orange roughy oil	40%
ouricury wax	50-55%
quinoa seed oil	6%
rye germ oil	11%
shark liver oil	60%
shea butter	9-13%
sperm whale oil	36%
sugar cane wax	18-80%
sunflower wax	25-45%
tall oil	9-23%
tall oil distillate	25-33%
Vegepure(TM) from wheat grains	70-90%
wheat germ oil	6%

“Substantivity” means the tendency of a material to resist being easily removed or the persistence of a treatment on the skin. For example, some sunscreen lotions are substantive because they form a film on the skin that is relatively water-insoluble. This, then, means that substantive materials resist removal or transfer by physical contact, sweating or washing.

- 5 Compositions of matter comprising waxes, oils and/or fats (lipids) containing at least 6% by weight unsaponifiable ingredients and at least 10% by weight saponifiable ingredients are subjected to an alkaline hydrolysis reaction to produce a non-foaming, substantive composition with unique

surfactant properties that may be used as an active ingredient or as a carrier for application of other active ingredients, e.g., as a carrier base for application of cosmetic, pharmaceutical or other active ingredients. Commercially available bio-based extracts that have high unsaponifiables include, but are not limited to, candelilla wax, carubba wax, jojoba oil, lanolin, lecithin, and shea butter.

5 The lipid subjected to the process of the invention may be a raw product or it can also undergo various refining and/or modification steps beforehand. Examples of refining processes which may be mentioned are the conventional processes of chemical or physical refining or the more specialized processes for the refining of shea butter, which make it possible in particular to retain or concentrate the maximum amount of unsaponifiable materials, thereafter subjecting such treated
10 materials to the process of the present invention.

The chemical refining which is preferentially used, being applied to the vegetable fats before they are subjected to the process according to the present invention, may be any conventional chemical refining process, in particular any process comprising the following steps:

Step 1: degumming involving insolubilization of the phosphatides with water, generally in
15 the presence of acid, most frequently phosphoric acid, and separation by decantation or centrifugation (continuous process);

Step 2: neutralization of the free fatty acids in the oil by the addition of a sodium hydroxide solution and separation of the soaps formed (called soap stock), most frequently by centrifugation followed by several washes with water, steps 1 and 2 often being
20 performed simultaneously in a continuous process;

Step 3: decolorization with activated bleaching clays at about 100°C under reduced vacuum, and filtration;

Step 4: deodorization operation necessary for removing the compounds responsible for the odors and flavors of an oil and for producing refined oil. This operation is carried
25 out in an apparatus called a deodorizer, the procedure involving heating of the oil to a high temperature (180°-220°C.) under a vacuum of the order of 4 torr (i.e. about 532 Pa) and a massive injection of steam to strip away impurities.

An alternate physical refining method is understood as a variant of the chemical refining process explained above, the difference being that the neutralization step with sodium hydroxide is
30 not performed and that the removal of the free fatty acids from the oil is effected during the deodorizing step. The refinement conditions selected during this physical refining method may require modification in order to retain the desired properties of the high unsaponifiables selected for use during the procedure for preparation of the present invention.

The extracts used as starting materials for the hydrolysis reaction according to the method

of the present invention may be in their raw or refined states. The extracts may also be alkoxylated, polymerized, acetylated, oxidized, reduced, concentrated, hydrogenated, partial hydrogenated, interesterified, double bond modified, randomized, refined, or otherwise modified before the hydrolysis reaction. Since many lipids have low concentrations or fractions (for example 1% or less
5 as discussed above) of unsaponifiables, the present invention encompasses the concentration of low fraction unsaponifiables into higher fractions, i.e., greater than 6%.

The products from the hydrolysis reaction of organic materials that produce unsaponifiables comprises a mixture of: a) polar hydrophilic salts (saponifiables); and b) non-polar, lipophilic materials (unsaponifiables), with the possibility of other materials also present, depending on the
10 source, state and form of the initial reactant.

The composition of materials created by the method according to the present invention are produced by the reaction of aqueous alkali metal hydroxides, e.g., NaOH, LiOH, KOH (the preferred hydroxide), CaOH, MgOH, and the like, with organic lipid compositions, usually plant extracts, oils, fats, or waxes (of the extracts or derivatives of the extracts) where the organic compositions contain
15 a high proportion of unsaponifiable materials (greater than 6%), and preferably as long chain esters.

Joboba oil may be examined as an example case. Refined jojoba oil contains various proportions of long chain diunsaturated esters. Hydrolysates of refined jojoba oil are nearly a 55:45 mixture of polar hydrophilic long chain salts (alkali salts) and relatively non-polar lipophilic materials (fatty alcohols). The lipophilic fraction is the unsaponifiable materials according to the
20 definition used in this document. The carbon chain lengths of both of these jojoba Hydrolysates include and vary from C₁₈ to C₂₄ and have ω-9 double bonds as part of each molecule. It has been found that the combination of saponifiable and unsaponifiable fractions of the Hydrolysates according to the present invention has properties that aid in the formulation of cosmetic, pharmaceutical, and other compositions.

The products that result from the hydrolysis of the lipids containing high percentages of unsaponifiable materials, as created during the practice of the present invention, whether used neat, blended, dissolved, dispersed, or emulsified with excipients, solvents, or carriers, can contain and impart useful properties to applied surfaces. These surfaces may be animate surfaces, particularly human skin, plant surfaces, and even the surfaces of inanimate objects, for example objects of wood,
30 fiber, or plastic. The properties can include, but are not limited to, substantivity, emulsification, hydration, and the like.

One of the above-mentioned properties, substantivity, is particularly useful in the field of lipstick, shampoos, conditioners, hair sheens, repellants, attractants, cosmetics, pharmaceuticals, and sunscreens. The property of substantivity is especially beneficial to hair care products, such as

“leave in” hair conditioners, where naturally derivatized materials that display substantivity are particularly commercially desirable. Substantivity is also particularly useful with sunscreen, sun block, or tanning formulations, as well as with insect repellants, such as tick, flea and fly repellants, and pesticides. Substantivity may also be beneficial when used on inanimate objects, such as with
5 air fresheners, antibacterial, antimildew, and antifungal agents, flystrips, pesticides, insecticides, insect repellants, herbicides, and the like.

It is theorized that the inclusion of the high levels of unsaponifiable materials in the organic material enables the Hydrolysates according to the present invention to display their unique combination of properties. The precise nature of the unsaponifiable materials within the oils, waxes,
10 fats or other natural extracts is not particularly important (except when a specific property is desired), and each of the variously available natural starting materials may differ significantly in their composition and types of unsaponifiables. For example, Jurinea extracts (e.g., the petroleum ether extracts of Jurinea) may comprise 40% by weight of pentacyclic triterpene alcohols together with their esters (myristate, palmitate, and acetate) as well as α -amyrin, β -amyrin, lupeol, and taraxasterol
15 such as ι -taraxasterol (*Lipids*, K. L. Mikolajczak et al., 1967, Vol. 2, No. 2, pp. 127-132). Briza oil may contain 20% by weight of lipids that are semi-solid, the lipid comprising 49% unsaponifiable digalactosylglycerides, 29% unsaponifiable monogalactosylglycerides and small amounts of conventional saponifiable triglycerides. The predominant fatty acids in the above oils are palmitic acid, oleic acid and linoleic acid (*Lipids*, C. R. Smith, Jr. et al., March 1966, Vol. 1, No. 2, pp. 123-
20 127).

The composition according to the present invention is preferably produced in a batch process using a large steam kettle equipped with a propeller mixer.

A measured quantity of potassium hydroxide pellets are added into the steam kettle with a measured quantity of distilled, deionized, or reverse osmosis purified water. The amount of
25 potassium hydroxide employed in order to completely saponify the free organic acid and/or organic acid ester can accordingly be calculated from the Saponification Value of the starting material and will, in theory, be the stoichiometric amount. In practice, however, it is preferred to employ slightly less than the stoichiometric amount of potassium hydroxide in order to ensure that the Hydrolysates that are formed are not contaminated with unused alkali. The amount of potassium hydroxide
30 employed can be considerably less than the stoichiometric amount, for example, as little as 50% of the stoichiometric amount or less may be used depending upon the desired result. It is to be understood, however, that an amount of potassium hydroxide in excess of the stoichiometric amount, for example, up to 10% more than the stoichiometric amount, can be employed if complete saponification of the organic acid or ester is to be achieved. Excess potassium hydroxide remaining

at the end of the reaction may be removed by traditional methods.

The potassium hydroxide pellets and water are stirred together with the propeller mixer until the potassium hydroxide pellets are dissolved. It is important to note, for safety purposes, that heat is generated during this step and the mixture is quite caustic. Individuals nearby should wear gloves,
5 eye and face protection, and clothing protection to avoid burns, both thermal and chemical.

Next, a measured quantity of a refined or derivatized organic material containing a high proportion of unsaponifiables, such as jojoba oil, is gently added to the steam kettle, taking care not to splash the caustic solution contained therein.

The steam kettle is heated to 90-95°C and held at that temperature range under constant
10 agitation for two hours. At this point, the resultant mixture should be pH tested. If the solution pH is greater than 10.0, continue heating the mixture under constant agitation at 90-95°C. Retest the solution periodically until the pH is 10.0 or less.

Once the pH is 10.0, or less, withdraw a sample for analysis. This sample should be analyzed by such methods as chromatography or by another like or similar method, to show that the
15 reaction has proceeded as desired.

The resultant Hydrolysate may then be diluted by adding a second measure quantity of water, or other diluent, to the steam kettle and stirred with the mixing propeller. Heat should be continuously applied, less than 80°C, until the mixture is homogeneous.

Once homogeneous, the Hydrolysate mixture is cooled to 60°C while continuing the mixing
20 with the propeller. The Hydrolysate mixture may then be transferred to a holding container and allowed to cool to room temperature before sealing the holding container.

Emulsification is the process of dispersing one material throughout another in separate droplets and, for industry's purposes, effecting a dispersion that will retain its physical characteristics for a period of one to two years at least. The influence on emulsifier type selected for use is related
25 to the ratio of hydrophilic and lipophilic character expressed by the emulsifier with reference to a similar, although reciprocal, character of the oil being emulsified. These two properties have been termed Hydrophilic-Lipophilic Balance (HLB) of the emulsifier and Required HLB of the oil. The HLB system is helpful to the emulsion formulator for the purpose of matching the appropriate emulsifier to a given oil. This matching is usually done experimentally, however, when the HLB
30 of an emulsifier and the HLB requirement of a given oil is known, this experimentation can be greatly reduced. The HLB of the present invention exhibits a unique property of being 3 to 4 HLB numbers wide and in the hydrophilic range. An emulsifier with a wide HLB effective range is advantageous due to the flexibility inherently imparted by such an emulsifier. The wide HLB effective range of the present invention also provides formulations with an extra margin for dealing

with unusual conditions such as pH, heat, cold, and the like, that may be encountered in the normal distribution of cosmetics, pharmaceutical and other bioactive products.

It was noted during an experiment that when a concentrated fly repellant (Purina Horse Spray Concentrate Insecticide) was diluted according to instructions the resulting mixture separated and required reintegration by shaking before use. This separation of components was eliminated by addition of the Hydrolysate according to the present invention, thus demonstrating the unique emulsification property of the Hydrolysate.

INDUSTRIAL APPLICABILITY

Below are described several example uses found for the Hydrolysates according to the present invention.

Example 1 Enhanced Skin Hydration

A Nova Meter is an impedance measuring device that is designed and commonly used to provide a non-invasive, objectively reproducible method of measurement for quantifying a biophysical character relative to hydration of the skin. Ten panelists participated in a skin hydration study that utilized a Nova Meter to register and record results. The test was conducted according to the following procedure.

A commercially available skin lotion was purchased and divided equally. Half was used as a control and half was used as a base into which 5% of a jojoba Hydrolysate was incorporated. The jojoba Hydrolysate was prepared according to the method disclosed in this invention. A baseline skin hydration reading was taken with the Nova Meter for each panelist in advance of any lotion application. The control and Hydrolysate containing lotions were applied to different areas of each panelist forearms. The Hydrolysate containing lotion was applied to the right forearm and the control lotion was applied to the left forearm. The Nova Meter was used to take skin hydration readings of the forearm areas to which each participant had applied each lotion. Multiple skin hydration readings were taken and recorded at two-hour intervals after lotion application. The results are illustrated in Figure 1.

The experiment resulted in a dramatic increase in skin hydration for most all test subjects in the test areas where the Hydrolysate formulation was applied, compared to the test areas of the control formulation. In general, 6 to 10 hours after application, the Hydrolysate lotion formulation demonstrated a 20% to 54% improvement in skin hydration over baseline areas. The Hydrolysate formulation showed a 10% to 47 % improvement in skin hydration over skin treated with the control formulation.

Example 2 – Reduces dehydration

Two make-up formulas were prepared: a Hydrolysate formulation containing 5% of a

Hydrolysate according to the present invention and a control formulation containing an extra 5% water. The 5% water was added to the control formulation to keep the remaining ingredient compositions the same between the two formulations. The control formulation was applied on the left forearm and the Hydrolysate formulation was applied on the right forearm.

5 A Nova Meter was used to take baseline hydration readings of each participant before make-up application and to take hydration readings at intervals of four and seven hours after application of each formulation. The results were averaged for each person using the control and Hydrolysate containing lotions to determine the percent difference in skin hydration from the baseline. The results are shown in Figure 2.

10 At four hours after make-up application, the average Nova Meter readings of the participants showed an increase in skin hydration of approximately 5% over baseline on areas to which the Hydrolysate formulation had been applied. A reduction in skin hydration of approximately 4% from baseline was observed on the areas with the control formulation. The difference between the Hydrolysate and control formulations was approximately 9%, with the Hydrolysate formulation
15 showing better hydration properties. In fact, the control formulation showed skin dehydration, which is not unusual for highly pigmented cosmetic formulations such as make-up and lipstick.

At seven hours after application, the average Nova Meter readings of the participants showed a reduction in skin hydration of approximately 4% below baseline on the areas with the Hydrolysate formulation. A reduction in skin hydration of approximately 6% below baseline was observed on
20 the areas with the control formulation. The hydration difference between the two make up formulations after seven hours was approximately 2%, with the Hydrolysate formulation continuing to show better hydration properties than the control formulation. Seven hours were required for the Hydrolysate makeup formulation to approach the drying level to the skin as compared to the control make up formulation.

25 Therefore, the incorporation of the Hydrolysates according to the present invention into typically drying make-up formations shows improved skin hydration properties compared to formulations not containing the Hydrolysates. In fact, the Hydrolysate formulation appears to hydrate the skin initially, as opposed to the dehydrating effect seen in the control make up formulation.

30 Example 3 – Enhanced Performance/Substantivity

Four different products for the treatment of fly abatement with animals, such as horses, were obtained. (Ceratex, Gnat-Away, Solitude, and Aqua-Pel.) Concentrated versions of these products were not available; therefore commercially available dilutions were used.

To each sample, either water or the Hydrolysate according to the present invention was

added, to make a 10% Hydrolysate containing solution. All formulations were thoroughly mixed with a stirrer until homogeneous. All formulations were transferred into spray bottles.

Four horses were selected to participate. The left side of each horse was sprayed with the control formulation. The right side of each horse was sprayed with the Hydrolysate formulation.

5 For eight (8) days, the number of flies on each horse's leg prior to re-application of any formulation was determined. With each of the four Hydrolysate formulations, the cumulative effect after eight days demonstrated a significant decrease in fly count. Figure 3-7 clearly shows that the Hydrolysate formulation produces a greater decrease in fly count than the control formulation. Thus, the inclusion of the Hydrolysate according to the present invention improves the cumulative performance of the
10 active materials transferred with the Hydrolysate in the commercially available fly abatement products.

The preferred embodiment(s) of the invention is described above in the Detailed Description of the Invention. While these descriptions directly describe the above embodiments, it is understood that those skilled in the art may conceive modifications and/or variations to the specific
15 embodiments shown and described herein. Any such modifications or variations that fall within the purview of this description are intended to be included therein as well. Unless specifically noted, it is the intention of the inventor that the words and phrases in the specification and claims be given the ordinary and accustomed meanings to those of ordinary skill in the applicable art(s). The foregoing description of a preferred embodiment and best mode of the invention known to the
20 applicant at the time of filing the application has been presented and is intended for the purposes of illustration and description. It is not intended to be exhaustive or to limit the invention to the precise form disclosed, and many modifications and variations are possible in the light of the above teachings. The embodiment was chosen and described in order to best explain the principles of the invention and its practical application and to enable others skilled in the art to best utilize the
25 invention in various embodiments and with various modifications as are suited to the particular use contemplated.

WHAT IS CLAIMED IS

- 1 1. A composition for topical application comprising products of hydrolysis of organic materials,
2 said organic materials further comprising at least 6 weight percent unsaponifiable materials.
- 1 2. The composition of claim 1 wherein said organic materials were pre-treated prior to
2 hydrolysis, said pre-treatment selected from the group of treatments consisting of
3 alkoxylation, polymerization, acetylation, oxidation, reduction, concentration, hydrogenation,
4 partial hydrogenation, interesterification, double bond modification, randomization, and
5 refinement.
- 1 3. The composition of claim 2 wherein said pre-treated organic materials further comprises
2 extracts selected from the group consisting of amaranth seed oil, anise seed oil, avocado seed
3 oil, barley oil, briza oil, buck wheat oil, candelilla wax, carnuba wax, cassia occidentalis oil,
4 coffee bean oil, deoiled lecithin, dog fish oil, esparto wax, oils from fungi and other
5 microorganisms, guayule plant extract, jojoba oil, jurinea oil, lanolin, laurel berry oil, olestra
6 (olean), olive oil concentrate (phytosqualene), olive seed oil, orange roughy oil, ouricury
7 wax, quinoa seed oil, rye germ oil, shark liver oil, shea butter, sperm whale oil, sugar cane
8 wax, sunflower wax, tall oil, tall oil distillate, Vegepure from wheat grains, and wheat germ
9 oil.
- 1 4. The composition of claim 1 comprising at least 20% by weight of unsaponifiables.
- 1 5. The composition of claim 2 comprising at least 20% by weight of unsaponifiables.
- 1 6. A substantive composition comprising the composition of claim 1 in combination with at

2 least one ingredient selected from the group consisting of emollients, conditioners, pigments,
3 dyes, pharmaceuticals, ultraviolet radiation absorbers, physical radiation blocks, insect
4 repellants, animal repellants, insecticides, pesticides, herbicides, animal attractants,
5 fragrances, and hormones.

1 7. A substantive composition comprising the composition of claim 2 in combination with at
2 least one ingredient selected from the group consisting of emollients, conditioners, pigments,
3 dyes, pharmaceuticals, ultraviolet radiation absorbers, physical radiation blocks, insect
4 repellants, animal repellants, insecticides, pesticides, herbicides, animal attractants,
5 fragrances, and hormones.

1 8. A substantive composition comprising the composition of claim 4 in combination with at
2 least one ingredient selected from the group consisting of emollients, conditioners, pigments,
3 dyes, pharmaceuticals, ultraviolet radiation absorbers, physical radiation blocks, insect
4 repellants, animal repellants, insecticides, pesticides, herbicides, animal attractants,
5 fragrances, and hormones.

1 9. A substantive composition comprising the composition of claim 5 in combination with at
2 least one ingredient selected from the group consisting of emollients, conditioners, pigments,
3 dyes, pharmaceuticals, ultraviolet radiation absorbers, physical radiation blocks, insect
4 repellants, animal repellants, insecticides, pesticides, herbicides, animal attractants,
5 fragrances, and hormones.

1 10. A method of providing substantive benefits to an animal subject comprising applying the
2 composition of claim 1 to the hair, skin, scales, or feathers of an animal subject.

- 1 11. A method of providing substantive benefits to an animal subject comprising applying the
2 composition of claim 2 to the hair, skin, scales, or feathers of an animal subject.
- 1 12. A method of providing substantive benefits to an animal subject comprising applying the
2 composition of claim 6 to the hair, skin, scales, or feathers of an animal subject.
- 1 13. A method of providing substantive benefits to an animal subject comprising applying the
2 composition of claim 7 to the hair, skin, scales, or feathers of an animal subject.
- 1 14. A method of providing substantive benefits to a botanical subject comprising applying the
2 composition of claim 1 to the surface of the botanical subject.
- 1 15. A method of providing substantive benefits to a botanical subject comprising applying the
2 composition of claim 2 to the surface of the botanical subject.
- 1 16. A method of providing substantive benefits to a botanical subject comprising applying the
2 composition of claim 6 to the surface of the botanical subject.
- 1 17. A method of providing substantive benefits to a botanical subject comprising applying the
2 composition of claim 7 to the surface of the botanical subject.
- 1 18. A method of providing substantive benefits to an inanimate object comprising applying the
2 composition of claim 1 to the surface of the inanimate object.
- 1 19. A method of providing substantive benefits to an inanimate object comprising applying the
2 composition of claim 2 to the surface of the inanimate object.

- 1 20. A method of providing substantive benefits to an inanimate object comprising applying the
2 composition of claim 6 to the surface of the inanimate object.
- 1 21. A method of providing substantive benefits to an inanimate object comprising applying the
2 composition of claim 7 to the surface of the inanimate object.
- 1 22. A composition for topical application comprising products of hydrolysis of organic materials,
2 said organic materials further comprising at least 6 weight percent long carbon chain
3 materials, said long carbon chain materials having at least 18 carbons in length.
- 1 23. The composition of claim 22 wherein said organic materials were pre-treated prior to
2 hydrolysis, said pre-treatment selected from the group of treatments consisting of
3 alkoxylation, polymerization, acetylation, oxidation, reduction, concentration, hydrogenation,
4 partial hydrogenation, interesterification, double bond modification, randomization, and
5 refinement.
- 1 24. The composition of claim 23 wherein said pre-treated organic materials further comprises
2 extracts selected from the group consisting of amaranth seed oil, anise seed oil, avocado seed
3 oil, barley oil, briza oil, buck wheat oil, candelilla wax, carnuba wax, cassia occidentalis oil,
4 coffee bean oil, deoiled lecithin, dog fish oil, esparto wax, oils from fungi and other
5 microorganisms, guayule plant extract, jojoba oil, jurinea oil, lanolin, laurel berry oil, olestra
6 (olean), olive oil concentrate (phytosqualene), olive seed oil, orange roughy oil, ouricury
7 wax, quinoa seed oil, rye germ oil, shark liver oil, shea butter, sperm whale oil, sugar cane
8 wax, sunflower wax, tall oil, tall oil distillate, Vegepure from wheat grains, and wheat germ
9 oil.

- 1 25. The composition of claim 22 comprising at least 20% by weight of unsaponifiables.
- 1 26. The composition of claim 23 comprising at least 20% by weight of unsaponifiables.
- 1 27. A substantive composition comprising the composition of claim 22 in combination with at
2 least one ingredient selected from the group consisting of emollients, conditioners, pigments,
3 dyes, pharmaceuticals, ultraviolet radiation absorbers, physical radiation blocks, insect
4 repellants, animal repellants, insecticides, pesticides, herbicides, animal attractants,
5 fragrances, and hormones.
- 1 28. A substantive composition comprising the composition of claim 23 in combination with at
2 least one ingredient selected from the group consisting of emollients, conditioners, pigments,
3 dyes, pharmaceuticals, ultraviolet radiation absorbers, physical radiation blocks, insect
4 repellants, animal repellants, insecticides, pesticides, herbicides, animal attractants,
5 fragrances, and hormones.
- 1 29. A substantive composition comprising the composition of claim 25 in combination with at
2 least one ingredient selected from the group consisting of emollients, conditioners, pigments,
3 dyes, pharmaceuticals, ultraviolet radiation absorbers, physical radiation blocks, insect
4 repellants, animal repellants, insecticides, pesticides, herbicides, animal attractants,
5 fragrances, and hormones.
- 1 30. A substantive composition comprising the composition of claim 26 in combination with at
2 least one ingredient selected from the group consisting of emollients, conditioners, pigments,
3 dyes, pharmaceuticals, ultraviolet radiation absorbers, physical radiation blocks, insect

4 repellants, animal repellants, insecticides, pesticides, herbicides, animal attractants,
5 fragrances, and hormones.

1 31. A method of providing substantive benefits to an animal subject comprising applying the
2 composition of claim 22 to the hair, skin, scales, or feathers of an animal subject.

1 32. A method of providing substantive benefits to an animal subject comprising applying the
2 composition of claim 23 to the hair, skin, scales, or feathers of an animal subject.

1 33. A method of providing substantive benefits to an animal subject comprising applying the
2 composition of claim 27 to the hair, skin, scales, or feathers of an animal subject.

1 34. A method of providing substantive benefits to an animal subject comprising applying the
2 composition of claim 28 to the hair, skin, scales, or feathers of an animal subject.

1 35. A method of providing substantive benefits to a botanical subject comprising applying the
2 composition of claim 22 to the surface of the botanical subject.

1 36. A method of providing substantive benefits to a botanical subject comprising applying the
2 composition of claim 23 to the surface of the botanical subject.

1 37. A method of providing substantive benefits to a botanical subject comprising applying the
2 composition of claim 27 to the surface of the botanical subject.

1 38. A method of providing substantive benefits to a botanical subject comprising applying the
2 composition of claim 28 to the surface of the botanical subject.

- 1 39. A method of providing substantive benefits to an inanimate object comprising applying the
2 composition of claim 22 to the surface of the inanimate object.
- 1 40. A method of providing substantive benefits to an inanimate object comprising applying the
2 composition of claim 23 to the surface of the inanimate object.
- 1 41. A method of providing substantive benefits to an inanimate object comprising applying the
2 composition of claim 27 to the surface of the inanimate object.
- 1 42. A method of providing substantive benefits to an inanimate object comprising applying the
2 composition of claim 28 to the surface of the inanimate object.
- 1 43. A sunscreen composition for topical application comprising physical radiation blocking
2 agents, radiation absorbing agents, and hydrolysis products of organic materials, said organic
3 materials further comprising at least 6 weight percent unsaponifiable materials.
- 1 44. The composition of claim 43 wherein said organic materials were pre-treated prior to
2 hydrolysis, said pre-treatment selected from the group of treatments consisting of
3 alkoxylation, polymerization, acetylation, oxidation, reduction, concentration, hydrogenation,
4 partial hydrogenation, interesterification, double bond modification, randomization, and
5 refinement.
- 1 45. The composition of claim 44 wherein said pre-treated organic materials further comprises
2 extracts selected from the group consisting of amaranth seed oil, anise seed oil, avocado seed
3 oil, barley oil, briza oil, buck wheat oil, candelilla wax, carnuba wax, cassia occidentalis oil,
4 coffee bean oil, deoiled lecithin, dog fish oil, esparto wax, oils from fungi and other

5 microorganisms, guayule plant extract, jojoba oil, jurinea oil, lanolin, laurel berry oil, olestra
6 (olean), olive oil concentrate (phytosqualene), olive seed oil, orange roughy oil, ouricury
7 wax, quinoa seed oil, rye germ oil, shark liver oil, shea butter, sperm whale oil, sugar cane
8 wax, sunflower wax, tall oil, tall oil distillate, Vegepure from wheat grains, and wheat germ
9 oil.

1 46. The composition of claim 43 comprising at least 20% by weight of unsaponifiables.

1 47. The composition of claim 44 comprising at least 20% by weight of unsaponifiables.

Fig. 1
Keri Lotion Nova Meter Study




Fig. 2 "Make-up on Arms" Nova Meter Study




Fig. 3 Average Percent Reduction




Fig. 4 Aquapel, Horse 1

Per Dr. McPhaul: Horse 1 is a dark brown horse, which means that he had to get closer to see the flies, which means that the flies didn't cooperate during the counting process.




Fig. 5 Ceratex, Horse 2

Per Dr. McPhaul: This horse was in need of shoeing, which means that dirt and manure caked up on the soles of his hooves, which caused him to have more flies congregated around his lower legs. He was constantly stomping the ground, so flies came and went with each stomp.




Fig. 6 GnatAway





Fig. 7 Solitude, Horse 4

Per Dr. McPhaul: This horse has a history of staying cleaner and seems to always have fewer flies on him, regardless of fly control methods.



INTERNATIONAL SEARCH REPORT

Inter al Application No

PCT/US 00/35469

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A61K7/48 A61K7/06

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

WPI Data, PAJ, EPO-Internal

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	FR 2 471 775 A (L'OREAL) 26 June 1981 (1981-06-26) the whole document	1,4,6,8, 10,12, 22,25, 27,29, 31,33, 43,46
X	US 5 759 555 A (LAWRENCE S. MOY) 2 June 1998 (1998-06-02) column 3, line 31 - line 47 column 3, line 63 -column 4, line 20 examples 1-4 claims 1,13 -/--	1,6,10, 12,22, 27,31,33

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents :

A document defining the general state of the art which is not considered to be of particular relevance

E earlier document but published on or after the international filing date

L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

O document referring to an oral disclosure, use, exhibition or other means

P document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

Z document member of the same patent family

Date of the actual completion of the international search

12 April 2001

Date of mailing of the international search report

20/04/2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Alvarez Alvarez, C

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 00/35469

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>WO 94 21764 A (DESLOG ET AL.) 29 September 1994 (1994-09-29)</p> <p>page 3, line 6 -page 8, line 14 examples 7-10 & US 5 679 393 A 21 October 1997 (1997-10-21) cited in the application -----</p>	<p>1-3, 6, 7, 10-13, 22-24, 27, 28, 31-34</p>

INTERNATIONAL SEARCH REPORT

information on patent family members

International Application No

PCT/US 00/35469

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
FR 2471775	A	26-06-1981	BE 886707 A	17-06-1981
			CA 1158168 A	06-12-1983
			CH 645267 A	28-09-1984
			DE 3047649 A	10-09-1981
			GB 2066071 A, B	08-07-1981
			IT 1141142 B	01-10-1986
			JP 1007961 B	10-02-1989
			JP 1525610 C	30-10-1989
			JP 56092805 A	27-07-1981
			US 4324802 A	13-04-1982
US 5759555	A	02-06-1998	US 5928659 A	27-07-1999
WO 9421764	A	29-09-1994	FR 2702773 A	23-09-1994
			AT 154825 T	15-07-1997
			DE 69403958 D	31-07-1997
			DE 69403958 T	22-01-1998
			DK 690904 T	02-02-1998
			EP 0690904 A	10-01-1996
			JP 9500916 T	28-01-1997
			US 5679393 A	21-10-1997